

Altria Altria Client Services

Abstract

Introduction: Few reports exist of exhaled breath levels of constituents during use of e-vapor products (EVPs). MarkTen[®] EVPs look similar to combustible cigarettes and consist of a battery and replaceable cartridges that contain propylene glycol (PG), glycerin (G), water, flavors and USP-grade tobacco-derived nicotine. The purpose of this study was to characterize the levels of selected constituents in exhaled breath during use of MarkTen[®] EVP.

Method: EVP users (n=32, 56% male) were randomized to use one of four flavor varieties each day for 4 days (2 menthol and 2 non-menthol flavors delivering 0.13 mg to 0.23 mg nicotine per machine measured 5 s puff). Subjects were instructed to take 10 puffs from each EVP using 5 s puff durations with 30 s inter puff intervals. The collection was performed once with a "sham" device (inactive battery + empty cartridge) and then again with the study EVP. Exhaled breath was collected after the first puff until 1 min after the last puff using a mouthpiece filter with a cryogenic condensation collection tube system. The collections were analyzed for PG, G, nicotine, menthol, formaldehyde, acrolein and acetaldehyde.

Results: Sham corrected exhaled breath levels (ANCOVA Least Square Means/10 puffs) across the flavor varieties ranged from: 1199.7 to 3354.5 µg for PG, 5366.8 to 6484.7 µg for G; 89.4 to 195.7 µg for nicotine, 0.17 (non-menthol product) to 31.01 µg (menthol product) for menthol, and from 0.25 to 0.34 µg for formaldehyde. Inter-subject variability was high for these analytes, with the largest range occurring for G where individual values ranged from 0 to 20,645 µg/10 puff sample. All acrolein and acetaldehyde exhaled breath measurements were below the detection limits. E-liquid consumption (cartridge weight change) during the 10 puff collections ranged from 34 to 41 mg across the four products. **Conclusions:** Detectable levels of PG, G, nicotine, menthol and formaldehyde are present in exhaled breath during use of EVPs. These values are currently being used in computational models to estimate room air levels under different room and use conditions.

Purpose

In Section VI (H) (2b) of the Draft Guidance for Premarket Tobacco Product Applications for Electronic Nicotine Delivery Systems (May 2016), the FDA recommends that applications:

"should provide data that adequately characterizes the likely impact of the new tobacco product on the health of both users and nonusers of tobacco products in order to support that marketing the new tobacco product would be appropriate for the protection of the public health."

The purpose of this study was to characterize the levels of selected constituents in the exhaled breath during use of the study EVPs. The results of this study will support our assessment of the likely impact of the EVPs on the health of non-users utilizing computational modeling to estimate the potential room air levels under different room and use conditions. The modeling results are presented at SRNT 2018, Poster 190 (Session 5) "Estimation of Second Hand Exposure Levels from ENDS and Conventional Cigarette Use Using Computational Modeling."

Objective

The primary objective of this study was to estimate the change in total amount of nicotine, glycerin, propylene glycol, menthol, formaldehyde, acetaldehyde, and acrolein in exhaled breath samples between sham (inactive battery and empty cartridge) and controlled product use for four EVPs.

e-Vapor Products

- MarkTen[®]A
- 2.5% nicotine by weight 4.0% nicotine by weight
- MarkTen[®] B
- MarkTen[®] D
- 3.5% nicotine by weight, Menthol
- 4.0% nicotine by weight, Menthol

Constituents Investigated

- Nicotine (N)
- Propylene glycol (PG)
- Formaldehyde(F)
- Acrolein (A)
- **Glycerin** (G) Menthol (M)
- Acetaldehyde (AA)


MarkTen[®] C

Study	Population	

Age	Mean	36.7
	Range	23 - 62
Gender	Male	18 (56%)
	Female	14 (44%)
Race	White or Caucasian	9 (28%)
	Black or African American	22 (69%)
	Other	1 (3%)
Current EVP type	Disposable / Cartridge based	11 (34%)
	Refillable	17 (53%)
	Disposable / Cartridge based + Refillable	4 (13%)
EVP use in the past week	Mean	6.5 days
	Range	4-7 days


Exhaled Breath Levels of Selected Constituents From Controlled Use of MarkTen[®] e-Vapor Products in Adult e-Vapor Users

Jeff Edmiston, Qiwei Liang, Sandra Miller, Jianmin Liu, Yuxi Zhao, Mohamadi Sarkar Altria Client Services LLC, Center for Research and Technology, 601 East Jackson Street, Richmond, VA 23219, USA SRNT 24th Annual Meeting, February 21 - 24, 2018, Baltimore, MD, USA




- response variable was sham-corrected analyte amount, study EVP, period and sequence were the fixed effects, sham sample analyte level was the covariate and subject nested within sequence was the random effect.

Figure 2. Exhaled Breath Sample Collection




Sham Sample Collections: 10 puffs of 5 second duration

Study EVP Sample Collections: 10 puffs of 5 second duration



Nicotine, PG, Glycerin, and Menthol



Collection #2 for: Formaldehyde, Acetaldehyde and Acrolein

EVP	Menthol (mg/puff) (SD)	Nicotine (mg/puff) (SD)	Glycerin (mg/puff) (SD)	Propylend Glycol (mg/puff) (S	Change (mg/puff)
MarkTen A	BLOQ	0.126 (0.006)	2.521 (0.123		
MarkTen B	BLOQ	0.201 (0.013)	2.885 (0.097) 1.011 (0.042	2) 5.275 (0.329)
MarkTen C	0.081 (0.003)	0.187 (0.005)	2.240 (0.075) 1.986 (0.070	D) 5.588 (0.177)
MarkTen D	0.130 (0.014)	0.232 (0.024)	2.294 (0.311) 1.940 (0.24	5) 5.972 (0.603)
EVP	Formaldel (µg/puff) (aldehyde ouff) (SD)	Acrolein µg/puff) (SD)	Cartridge Weight Change (µg/puff) (SD)
MarkTen A	0.100 (0.0	29) 0.06	65 (0.008)	BLOQ	5.662 (0.383)
MarkTen B	0.178 (0.0	17) 0.09	90 (0.008)	BLOQ	5.362 (0.314)
MarkTen C	0.187 (0.1	13) 0.07	75 (0.048)	BLOQ	5.603 (0.458)
MarkTen D	0.133 (0.0	23) 0.05	51 (0.006)	BLOQ	6.088 (0.372)

This poster may be accessed at www.altria.com/ALCS-Science

ANCOVA Least Square Means (95% CI) of Sham Corrected Exhaled Breath from 10 Controlled Puffs

			Propylene		
udy EVP	Nicotine	Glycerin	Glycol	Menthol	Formaldehyde
	[ug]	[ug]	[ug]	[ug]	[ug]
Α	89.44	5,972.3	1678.4	0.17	0.25
	(15.77, 163.11)	(4,191.1, 7,753.6)	(592.14, 2764.6)	(–2.92, 3.25)	(0.12, 0.38)
В	195.70	6,099.5	1199.7	0.35	0.25
	(122.01, 269.38)	(4,317.8, 7,881.2)	(113.74, 2285.7)	(–2.70, 3.40)	(0.12, 0.38)
C	168.83	6,484.7	3354.5	21.11	0.34
	(95.17, 242.50)	(4,701.7, 8,267.6)	(2266.6, 4442.3)	(18.06, 24.16)	(0.21, 0.47)
D	182.65	5,366.8	2511.0	31.01	0.30
	(108.63, 256.67)	(3,575.9, 7,157.7)	(1416.1, 3605.9)	(27.91,34.12)	(0.17, 0.43)

Estimated Mean e-Liquid Consumed* During Exhaled Breath Collections

Study E\/D		Α	В	С	D
Study EVP					
	Sample Size	32	32	32	32
Collection #1	Mean (mg)	35	34	40	37
(N, G, PG, M)	SD	8.7	11.4	12.6	12.3
Collection #2	Mean (mg)	36	36	41	38
(F, AA, A)	SD	11.5	11.9	14.6	12.4

*Cartridge weight change over the 10 puff session

Estimated Amount of Constituent Exhaled per mg e-Liquid Consumed*

Study EVP	Nicotine: μg exhaled / mg cartridge weight change	Glycerin: µg exhaled / mg cartridge weight change	PG: µg exhaled / mg cartridge weight change	Menthol: μg exhaled / mg cartridge weight change	Formaldehyde: µg exhaled / mg cartridge weight change	
Α	2.56	170.64	47.95	_	0.0069	
В	5.76	179.40	35.29	_	0.0069	
С	4.22	162.12	83.86	0.53	0.0083	
Ğ				0.00	0.0000	
_						
D	4.94	145.05	67.86	0.84	0.0079	
*Exhaled breath values from statistical model results and e-liquid consumption from cartridge weight change over the						

10 puff session

Conclusion

- The measured constituents in exhaled breath samples during study EVP use are highly variable.
- Nicotine, Glycerin, PG and Formaldehyde were detected in the exhaled breath for all four study EVPs. Menthol was detected in two menthol products.
- Acetaldehyde and Acrolein were below the detectable limit in exhaled breath during use of any of the four study EVPs. These values are currently being used in computational models to estimate room air levels under different room and use
- conditions (SRNT 2018, Poster 190, Session 5).