Evaluation Summary of Vanillin for Use as a Cigarette Ingredient

Vanillin is listed generally recognized as safe (GRAS) by United States Food and Drug Administration (FDA), and is also approved as a food additive (21 CFR § 182.60). Vanillin is designated as GRAS by the Flavor Extract Manufacturers Association (FEMA No. 3107)¹ and is approved for use in food by the Joint FAO/WHO Expert Committee on Food Additives (JECFA)^{2,3} and the Council of Europe (CoE No. 107).⁴

Vanillin is perhaps one of the most recognizable aroma chemicals in the world. Its desirable characteristic flavour and aroma have led to its widespread use in the food industry.⁵⁻¹⁰ In addition to its use in food, vanillin is used extensively as a fragrance in cosmetics, personal care products air fresheners and as a masking agent for otherwise odorous chemicals. It is also used in confectionery industries, and as a pharmaceutical excipient in oral solutions, tablet coatings, syrups and in powders to mask otherwise unpleasant tasting pharmaceutical preparations.¹¹

Toxicity studies have shown that vanillin is of low toxicity with lethal oral doses in rodents of approximately 1-1.5 g/kg.¹²⁻¹⁸ Short term and chronic studies have demonstrated the safety of daily exposure to vanillin in food.¹⁸⁻²¹ Teratology studies with rats have not demonstrated any danger to the fetus.²²⁻²⁶ Genotoxicity studies in bacterial and mammalian cell systems did not reveal any mutagenic properties at µM concentrations but mM concentrations induced sister chromatid exchanges in normal human lymphocytes.^{13,26-48} Vanillin has been reported in many studies to possess anticarcinogenic qualities.^{12,49-53} Evaluation of biochemical systems reveals that vanillin can suppress many of the enzymes that are required for the metabolic activation of known mutagens and carcinogens.⁵⁴⁻⁶⁶ *In vitro* studies with bacterial and mammalian cells demonstrated that, in the presence of some mutagens, vanillin would reduce their genotoxicity but that it can also promote the genotoxicity of others.^{26,42-47,66-69} Antimutagenic studies *in vivo* have described a similar dual effect of vanillin, which also proved to be dependent on the mutagen used.^{12,49-51}

Vanillin is currently used worldwide at levels below 500 ppm in selected cigarette brands manufactured and/or distributed by Philip Morris USA Inc. (PM USA) and/or Philip Morris Products SA (PMP SA). Vanillin is a natural component of tobacco (0.9-1.8 ppm).⁷⁰ Vanillin may be applied directly to the tobacco as an additive, flavoring or flavoring agent, and as such, may be subject to pyrolysis-type reactions during the smoking process. Vanillin may also be applied to the filter as a flavoring material where it would not be subjected to pyrolysis temperatures.

As suggested by purge and trap studies conducted by PM USA, vanillin would be expected to distill at 100°C prior to the burning cone of the tobacco.⁷¹ At the higher temperatures used in the pyrolysis studies conducted by PM USA, the results suggested that vanillin would not pyrolyze extensively during the smoking process and would be delivered to the smoke intact.⁷²

Vanillin was part of a PM USA testing program that was designed to evaluate the potential effects of 333 ingredients added to typical commercial blended test cigarettes on selected biological and chemical endpoints.⁷³⁻⁷⁶ Three pairs of test cigarettes were produced, each containing different groups of ingredients. Vanillin was added to two pairs at target levels of 23 ppm, 70 ppm, 276 ppm and 829 ppm. No significant effects were noted in cytotoxicity,

mutagenic studies or in respiratory tract endpoints in 90-day rat inhalation studies. In addition, smoke chemistry studies from cigarettes containing a mixture of flavors including vanillin did not significantly alter the smoke chemistry profile compared to control cigarettes. Based on the results of these studies, the authors concluded that these ingredients (including vanillin) added to tobacco do not add significantly to the overall toxicity of cigarettes.

Currently, information is only available for tests utilizing vanillin in a mixture of ingredients applied to cigarette tobacco. Studies are ongoing to address the use of vanillin as a single ingredient. Published studies show there is no meaningful difference in the composition or toxicity of smoke from cigarettes with added ingredients (including vanillin) compared to the smoke from cigarettes without added ingredients.⁷³⁻⁸¹ Based on the best available data, the ingredients used in PM USA and/or PMP SA cigarettes do not increase the overall toxicity of cigarette smoke.

References

- Hall, R.L. and Oser, B.L. (1965) Recent Progress in the Consideration of Flavoring Ingredients Under the Food Additives Amendment. III GRAS Substances. *Food Technology* 19(2):151-197.
- 2. Eleventh Report of the Joint FAO/WHO Expert Committee on Food Additives (1967) FAO Nutrition Meeting Report Series. Report No. 44, Wld Hlth Org. techn. Rep. Ser. 1968, 383.
- 3. Fifty-seventh Meeting of the Joint FAO/WHO Expert Committee on Food Additives (2001) TRS 909-JECFA 57/84.
- 4. CoE (2000) *Chemically-Defined Flavouring Substances*. Council of Europe Publishing, F-67075 Strasbourg Cedex, France. p.120.
- 5. Arctander, S. (1960) Perfume and flavor materials of natural origin. Unknown 2:638-647.
- 6. Belay, M.T. and Poole, C.F. (1993) Determination of vanillin and related flavor compounds in natural vanilla extracts and vanilla-flavored foods by thin layer chromatography and automated multiple development. *Chromatographia* 37(7-8):365-373.
- 7. Burdock, G.A. (1995) *Fenaroli's handbook of flavor ingredients*. Third Edition Edition. Vol. 1 & 2. CRC Press, Boca Raton, FL.
- 8. Burdock, G.A. (1997) *Encyclopedia of food and color additives*. Vol. 1, 2, and 3. CRC Press, Boca Raton, FL.
- 9. Clark, G.S. (1990) Vanillin. Perfumer and Flavorist 15:45-56.
- 10. Kahan, S. and Krueger, D.A. (1997) Liquid chromatographic method for determination of vanillin and ethyl vanillin in imitation vanilla extract (modification of AOAC official method 990.25): collaborative study. *Journal of AOAC International* 80(3):564-570.
- American Pharmaceutical Association (1994) Handbook of pharmaceutical excipients. 2nd edition Edition. (A. Wade and P. J. Weller). American Pharmaceutical Association, Washington, DC. p.651.
- 12. Inouye, T.; Sasaki, Y.F.; Imanishi, H.; Watanebe, M.; Ohta, T. and Shirasu, Y. (1988) Suppression of mitomycin C-induced micronuclei in mouse bone marrow cells by posttreatment with vanillin. *Mutat Res* 202(1):93-95.
- 13. Bonin, A.M. and Baker, R.S.U. (1980) Mutagenicity testing of some approved food additives with the Salmonella microsome assay. *Food Tech in Australia* 32(12):608-611.

- Taylor, J.M.; Jenner, P.M. and Jones, W.I. (1964) A comparison of the toxicity of some allyl, propenyl and propyl compounds in the rat. *Toxicology and Applied Pharmacology* 6:378-387.
- 15. Hodge, H.C. and Downs, W.L. (1961) The approximate oral toxicity in rats of selected household products. *Toxicology and Applied Pharmacology* 3:689-695.
- Jenner, P.M.; Hagan, E.C.; Taylor, J.M.; Cook, E.L. and Fitzhugh, O.G. (1964) Food flavourings and compounds of related structure. I. Acute oral toxicity. *Food and Cosmetics Toxicology* 2:327-343.
- Spector, W.S. (1956) Handbook of toxicology, Volume 1: Acute toxicities of solids, liquids and gases to laboratory animals. (W. S. Spector). W.B.Saunders Company, Philadelphia, PA. p.312-313.
- 18. Deichmann, W. and Kitzmiller, K.V. (1940) On the toxicity of vanillin and ethyl vanillin for rabbits and rats. *Journal of the American Pharmaceutical Association* 29(10):425-428.
- Hagan, E.C.; Hansen, W.H.; Fitzhugh, O.G.; Jenner, P.M.; Jones, W.I.; Taylor, J.M.; Long, E.L.; Nelson, A.A. and Brouwer, J.B. (1967) Food flavourings and compounds of related structure. II. Subacute and chronic toxicity. *Food Cosmet Toxicol* 5(2):141-157.
- 20. Gaworski, C.L.; Vollmuth, T.A.; Dozier, M.M.; Heck, J.D.; Dunn, L.T.; Ratajczak, H.V. and Thomas, P.T. (1994) An immunotoxicity assessment of food flavouring ingredients. *Food Chem Toxicol* 32(5):409-415.
- 21. Srinivasan, M.R. and Chandrasekhara, N. (1992) Comparative influence of vanillin & capsaicin on liver & blood lipids in the rat. *Indian J Med Res* 96:133-135.
- 22. Morita, Y.; Matsuda, H.; Mizutani, M. and Ishihara, Y. (1987) Effect of antimutagens on the teratogenicity of mitomycin C and alkylating agents. *Teratology* 36(3):467.
- 23. Morita, Y. and Mizutani, M. (1988) Effect of antimutagens on the teratogenicity of alkylating agents. *Teratology* 38(5):539.
- 24. Morita, Y. and Mizutani, M. (1988) Effects of antimutagens on the teratogenicity of nmethyl-n'-nitro-n-nitrosoguanidine in mice. 28:157-167. *Congenital Anomalies. (Senten Ijo)* 28:157-167.
- 25. Morita, Y. and Shiota, K. (1997) Effects of antimutagens on the teratogenicity of enu in mice. *Congenital Anomalies* 37(3):314.
- 26. Imanishi, H.; Sasaki, Y.F.; Matsumoto, K.; Watanabe, M.; Ohta, T.; Shirasu, Y. and Tutikawa, K. (1990) Suppression of 6-TG-resistant mutations in V79 cells and recessive spot formations in mice by vanillin. *Mutat Res* 243(2):151-158.
- Ishidate, M., Jr.; Sofuni, T.; Yoshikawa, K.; Hayashi, M.; Nohmi, T.; Sawada, M. and Matsuoka, A. (1984) Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 22(8):623-636.

- Mortelmans, K.; Haworth, S.; Lawlor, T.; Speck, W.; Tainer, B. and Zeiger, E. (1986) Salmonella mutagenicity tests. II. Results from the testing of 270 chemicals. Environmental Mutagenesis 8(Suppl 7):1-119.
- 29. Nagabhushan, M. and Bhide, S.V. (1985) Mutagenicity of chili extract and capsaicin in short-term tests. *Environ Mutagen*. 7(6):881-888.
- Takahashi, K.; Sekiguchi, M. and Kawazoe, Y. (1990) Effects of vanillin and o-vanillin on induction of DNA-repair networks: modulation of mutagenesis in Escherichia coli. *Mutat Res* 230(2):127-134.
- 31. Sivaswamy, S.N.; Balachandran B. and Balanehru S. (1991) South Indian foods: contaminants and their effects. *Bulletin of Environmental and Contamination Toxicology* 47:251-260.
- Watanabe, K.; Ohta, T.; Watanabe, M.; Kato, T. and Shirasu, Y. (1990) Inhibition of induction of adaptive response by o-vanillin in Escherichia coli B. *Mutat Res* 243(4):273-280.
- Watanabe, K.; Ohta, T.; Kato, T.; Watanabe, M. and Shirasu, Y. (1989) O-Vanillin enhances mutagenesis induced by MNNG in *Escherichia coli*. *Environ Mol Mutagen* 14(Suppl 15):215.
- 34. Watanabe, K.; Ohta, T. and Shirasu, Y. (1989) Enhancement and inhibition of mutation by ovanillin in Escherichia coli. *Mutat Res* 218(2):105-109.
- Ohta, T.; Watanabe, M.; Watanabe, K.; Shirasu, Y. and Kada, T. (1986) Inhibitory effects of flavourings on mutagenesis induced by chemicals in bacteria. *Food Chem Toxicol* 24(1):51-54.
- 36. Sato, T.; Ose, Y.; Nagase, H. and Hayase, K. (1987) Mechanism of the desmutagenic effect of humic acid. *Mutat Res* 176(2):199-204.
- 37. Fahrig, R. (1996) Anti-mutagenic agents are also co-recombinogenic and can be converted into co-mutagens. *Mutat Res* 350(1):59-67.
- 38. Jansson, T.; Curvall, M.; Hedin, A. and Enzell, C.R. (1986) In vitro studies of biological effects of cigarette smoke condensate. II. Induction of sister-chromatid exchanges in human lymphocytes by weakly acidic, semivolatile constituents. *Mutat Res* 169(3):129-139.
- 39. Jansson, T.; Curvall, M.; Hedin, A. and Enzell, C.R. (1988) In vitro studies of the biological effects of cigarette smoke condensate. III. Induction of SCE by some phenolic and related constituents derived from cigarette smoke. A study of structure-activity relationships. *Mutat Res* 206(1):17-24.
- 40. Jansson, T. and Zech, L. (1987) Effects of vanillin on sister-chromatid exchanges and chromosome aberrations in human lymphocytes. *Mutat Res* 190(3):221-224.

- 41. Kasamaki, A. and Urasawa, S. (1985) Transforming potency of flavoring agents in Chinese hamster cells. *J Toxicol Sci* 10(3):177-185.
- 42. Sanyal, R.; Darroudi, F.; Parzefall, W.; Nagao, M. and Knasmuller, S. (1997) Inhibition of the genotoxic effects of heterocyclic amines in human derived hepatoma cells by dietary bioantimutagens. *Mutagenesis* 12(4):297-303.
- 43. Sasaki, Y.F.; Imanishi, H.; Ohta, T. and Shirasu, Y. (1987) Effects of antimutagenic flavourings on SCEs induced by chemical mutagens in cultured Chinese hamster cells. *Mutat Res* 189(3):313-318.
- 44. Sasaki, Y.F.; Imanishi, H.; Ohta, T. and Shirasu, Y. (1987) Effects of vanillin on sisterchromatid exchanges and chromosome aberrations induced by mitomycin C in cultured Chinese hamster ovary cells. *Mutat Res* 191(3-4):193-200.
- 45. Tamai, K.; Tezuka, H. and Kuroda, Y. (1992) Different modifications by vanillin in cytotoxicity and genetic changes induced by EMS and H2O2 in cultured Chinese hamster cells. *Mutat Res* 268(2):231-237.
- 46. Matsumura, H.; Watanabe, K. and Ohta, T. (1993) o-Vanillin enhances chromosome aberrations induced by alkylating agents in cultured Chinese hamster cells. *Mutat Res* 298(3):163-168.
- Keshava, C.; Keshava, N.; Ong, T.M. and Nath, J. (1998) Protective effect of vanillin on radiation-induced micronuclei and chromosomal aberrations in V79 cells. *Mutat Res* 397(2):149-159.
- 48. Schlotzhauer, W.S.; Walters, D.B.; Snook, M.E. and Higman, H.C. (1978) Characterization of catechols, resorcinols, and hydroquinones in an acidic fraction of cigarette smoke condensate. *Journal of Agricultural and Food Chemistry* 26(6):1277-1281.
- 49. Inouye, T.; Sasaki, Y.F.; Imanishi, H.; Watanabe, M.; Kato, T.; Matsumoto, K.; Ohta, T. and Shirasu, Y. (1989) Suppressive effect of vanillin on micronuclei induced by mitomycin c or x-rays. *Environmental and Molecular Mutagenesis* 14(Suppl 15):93-94.
- 50. Sasaki, Y.F.; Ohta, T.; Imanishi, H.; Watanabe, M.; Matsumoto, K.; Kato, T. and Shirasu, Y. (1990) Suppressing effects of vanillin, cinnamaldehyde, and anisaldehyde on chromosome aberrations induced by X-rays in mice. *Mutat Res* 243(4):299-302.
- 51. Tsuda, H.; Uehara, N.; Iwahori, Y.; Asamoto, M.; Iigo, M.; Nagao, M.; Matsumoto, K.; Ito, M. and Hirono, I. (1994) Chemopreventive effects of beta-carotene, alpha-tocopherol and five naturally occurring antioxidants on initiation of hepatocarcinogenesis by 2-amino-3-methylimidazo[4,5-f]quinoline in the rat. *Jpn J Cancer Res* 85(12):1214-1219.
- 52. Akagi, K.; Hirose, M.; Hoshiya, T.; Mizoguchi, Y.; Ito, N. and Shirai, T. (1995) Modulating effects of ellagic acid, vanillin and quercetin in a rat medium term multi-organ carcinogenesis model. *Cancer Lett* 94(1):113-121.

- 53. Steele, V.E.; Moon, R.C.; Lubet, R.A.; Grubbs, C.J.; Reddy, B.S.; Wargovich, M.; McCormick, D.L.; Pereira, M.A.; Crowell, J.A.; Bagheri, D. and . (1994) Preclinical efficacy evaluation of potential chemopreventive agents in animal carcinogenesis models: methods and results from the NCI Chemoprevention Drug Development Program. *J Cell Biochem Suppl* 20:32-54.
- 54. Sammons, H.G. and Williams, R.T. (1941) Studies in detoxication. 12. The metabolism of vanillin and vanillic acid in the rabbit. The identification of glucurovanillin and the structure of glucurovanillic acid. *Biochemical Journal*. 35:1175-1189.
- 55. Wong, K.P. and Sourkes, T.L. (1966) Metabolism of vanillin and related substances in the rat. *Can. J Biochem* 44(5):635-644.
- 56. Strand, L.P. and Scheline, R.R. (1975) The metabolism of vanillin and isovanillin in the rat. *Xenobiotica* 5(1):49-63.
- 57. Scheline, R.R. (1972) The metabolism of some aromatic aldehydes and alcohols by the rat intestinal microflora. *Xenobiotica* 2(3):227-236.
- 58. Aboobaker, V.S.; Balgi, A.D. and Bhattacharya, R.K. (1994) In vivo effect of dietary factors on the molecular action of aflatoxin B1: role of non-nutrient phenolic compounds on the catalytic activity of liver fractions. *In Vivo* 8(6):1095-1098.
- 59. Cruickshank, D. and Sansom, L.N. (1993) Inhibition of rat liver aryl sulphotransferase by the food additive vanillin. *European Journal of Pharmaceutical Science* 1:41-47.
- 60. Bamforth, K.J.; Jones, A.L.; Roberts, R.C. and Coughtrie, M.W. (1993) Common food additives are potent inhibitors of human liver 17 alpha-ethinyloestradiol and dopamine sulphotransferases. *Biochem Pharmacol* 46(10):1713-1720.
- 61. Morse, M.A.; Kresty, L.A. and Toburen, A.L. (1995) Inhibition of metabolism of 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone by dietary benzaldehydes. *Cancer Lett* 97(2):255-261.
- 62. Kazakoff, K.; Iversen, P.; Lawson, T.; Baron, J.; Guengerich, F.P. and Pour, P.M. (1994) Involvement of cytochrome P450 2E1-like isoform in the activation of N-nitrosobis(2oxopropyl)amine in the rat nasal mucosa. *Eur J Cancer B Oral Oncol.* 30B(3):179-185.
- 63. Thompson, D.C. and Trush, M.A. (1986) The toxicological implications of the interaction of butylated hydroxytoluene with other antioxidants and phenolic chemicals. *Food Chem Toxicol* 24(10-11):1189-1195.
- 64. Sawa, T.; Nakao, M.; Akaike, T.; Ono, K. and Maeda, H. (1999) Alkylperoxyl radicalscavenging activity of various flavonoids and other phenolic compounds: implications for the anti-tumor-promoter effect of vegetables. *J Agric Food Chem* 47(2):397-402.
- 65. Laughton, M.J.; Evans, P.J.; Moroney, M.A.; Hoult, J.R. and Halliwell, B. (1991) Inhibition of mammalian 5-lipoxygenase and cyclo-oxygenase by flavonoids and phenolic dietary

additives. Relationship to antioxidant activity and to iron ion-reducing ability. *Biochem Pharmacol* 42(9):1673-1681.

- 66. Sharma, S.; Stutzman, J.D.; Kelloff, G.J. and Steele, V.E. (1994) Screening of potential chemopreventive agents using biochemical markers of carcinogenesis. *Cancer Res* 54(22):5848-5855.
- 67. Arnold, J.T.; Wilkinson, B.P.; Sharma, S. and Steele, V.E. (1995) Evaluation of chemopreventive agents in different mechanistic classes using a rat tracheal epithelial cell culture transformation assay. *Cancer Res* 55(3):537-543.
- 68. Keshava, C.; Keshava, N.; Whong, W.Z.; Nath, J. and Ong, T.M. (1997) Inhibition of methotrexate-induced chromosomal damage by vanillin and chlorophyllin in V79 cells. *Teratog. Carcinog. Mutagen.* 17(6):313-326.
- 69. Tamai, K.; Tezuka, H.; Kuroda, Y. and Kada, T. (1988) Studies on the combined effects of environmental mutagens on mammalian cells in culture. *Mutat Res* 203:391.
- 70. Demian, B.A. (1993) Trace analysis of vanillin in tobacco. *Journal of Liquid Chromatography* 16(16):3563-3574.
- 71. PM USA (2001) P&T/GC/MS Analysis of Vanillin. Request 20010503. Scan TF131RFD. Unpublished Internal Report.
- 72. PM USA (2001) Pyrolysis GC/MS Analysis of Vanillin. Request 20010503. Scan P010503. Unpublished Internal Report.
- 73. Carmines, E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 1: Cigarette design, testing approach, and review of results. *Food and Chemical Toxicology* 40:77-91.
- 74. Roemer, E.; Tewes, F.J.; Meisgen, T.J.; Veltel, D. and Carmines, E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 3: *In vitro* genotoxicity and cytotoxicity. *Food and Chemical Toxicology* 40:105-111.
- 75. Rustemeier, K.; Stabbert, R.; Haussmann, H.J.; Roemer, E. and Carmines E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 2: Chemical composition of mainstream smoke. *Food and Chemical Toxicology* 40:93-104.
- 76. Vanscheeuwijck, P.M.; Teredesai, A.; Terpstra, P.M.; Verbeek, J.; Kuhl, P.; Gerstenberg, B.; Gebel, S. and Carmines E.L. (2002) Evaluation of the potential effects of ingredients added to cigarettes. Part 4: Subchronic inhalation toxicity. *Food and Chemical Toxicology* 40:113-131.
- 77. Gaworski, C.L.; Dozier, M.M.; Heck, J.D.; Gerhart, J.M.; Rajendran, N.; David, R.M.; Brennecke, L.H. and Morrissey, R. (1998) Toxicologic evaluation of flavor ingredients added to cigarette tobacco: 13 week inhalation exposures in rats. *Inhal. Toxicol.* 10:357-381.

- 78. Gaworski, C.L.; Heck, J.D.; Bennett, M.B. and Wenk, M.L. (1999) Toxicologic evaluation of flavor ingredients added to cigarette tobacco: skin painting bioassay of cigarette smoke condensate in SENCAR mice. *Toxicology* 139(1-2):1-17.
- 79. Doull, J.; Frawley, J.P.; George, W.J.; Loomis, T.A.; Squire, R.A. and Taylor, S.L. (1994) A safety assessment of ingredients added to tobacco in the manufacturing of cigarettes. Covington and Burling, Washington, D.C.
- 80. Doull, J.; Frawley, J.P.; George, W.J.; Loomis, T.A.; Squire, R.A. and Taylor, S.L. (1998) A safety assessment of ingredients added to tobacco in the manufacturing of cigarettes. Covington and Burling, Washington, D.C.
- Baker, R.R.; Massey, E.D. and Smith, G. (2004) An overview of the effects of tobacco ingredients on smoke chemistry and toxicity. *Food and Chemical Toxicology* 42S(Supplement 1):S53-S83.