

For Bones. Skip Longitudinal and Axial if Multifocal is selected.

Longitudinal Location in Bone (Lesion Center)

- | | |
|-------------------------------------|---|
| ☐ Epiphysis | ☐ Diaphysis |
| ☐ Metaphysis | ☐ Not Applicable |

Axial Location

- | | |
|--|---|
| ☐ Central medullary | ☐ Intracortical |
| ☐ Eccentric medullary | ☐ Surface lesion |

Pattern of Bone Involvement

- ☐ Aggressive
- ☐ Nonaggressive

☐ Cortical Destruction

☐ Soft Tissue Mass/Extension/Component

☐ Surrounding Edema

Matrix Mineralization/Other Matrix

- | | |
|------------------------------------|---|
| ☐ Chondroid | ☐ Both/Indeterminate |
| ☐ Osteoid | ☐ Other |

Periosteal Reaction

- | | |
|-------------------------------------|--|
| ☐ Aggressive | ☐ Nonaggressive |
|-------------------------------------|--|

☐ Joint Involvement

☐ Neurovascular Involvement

☐ Fluid Levels

Is biopsy needed? ☐ Yes

Is the lesion more likely benign or malignant?

- ☐ Benign
- ☐ Malignant

Most Likely Diagnosis

- ☐ Adamantinoma
- ☐ Aneurysmal Bone cyst
- ☐ Chondroblastoma
- ☐ Chondrosarcoma
- ☐ Enchondroma
- ☐ Enchondromatosis
- ☐ Enchondromatosis to sarcoma
- ☐ Ewing Sarcoma
- ☐ Fibrosarcoma/MFH
- ☐ Fibrous Dysplasia
- ☐ Giant Cell Tumor
- ☐ Hemangioma
- ☐ Hereditary Multiple Exostoses
- ☐ Langerhans Cell Histiocytosis
- ☐ Lipoma
- ☐ Lymphoma
- ☐ Metastasis
- ☐ Multiple Myeloma/Plasmacytoma
- ☐ Nonossifying fibroma
- ☐ Osteoblastoma
- ☐ Osteochondroma to Sarcoma (single or multiple)
- ☐ Osteochondroma, Solitary
- ☐ Osteoid Osteoma
- ☐ Osteosarcoma, Intramedullary
- ☐ Osteosarcoma, Surface
- ☐ Unicameral Bone Cyst