

Children's Wisconsin Clinical Nutrition Department



PICU: Pre-Test- Answers

1. When is the best time to start enteral feeds if medically appropriate?
 - a. Within 6 hours of admission
 - b. Within 24 hours of admission**
 - c. Within 72 hours of admission
 - d. Within 7 days of admission
2. Total fluid goal (including nutrition support) for an intubated patient should usually be ____% maintenance fluid need
 - a. 50%
 - b. 80%**
 - c. 100%
 - d. 120%
3. True or **False**: If a well-nourished 8 year old child is not appropriate to start enteral nutrition on day 3 of admission, should start parenteral nutrition.
4. Your patient has been NPO for 4 days and is now on BiPap 24 hours/day due to respiratory status, what is the best nutrition intervention to recommend?
 - a. Remain NPO
 - b. Start Clear Liquid diet
 - c. Start bolus NG feeds
 - d. Start NJ feeds**
5. Which forms of dialysis should a renal enteral formula be recommended? (choose all that apply)
 - a. Hemodialysis**
 - b. Peritoneal Dialysis**
 - c. Continuous Renal Replacement Therapy/CVVH
6. Which micronutrients should be recommended for Burns >20% TBSA?
 - a. Zinc, Ascorbic Acid, Iron, Calcium
 - b. Ascorbic Acid, Vitamin D, Iron, Copper
 - c. Zinc, Ascorbic Acid, Vitamin A, Vitamin D**
 - d. Vitamin A, Vitamin D, Vitamin E, Magnesium