



Australian Government
**Australian Pesticides and
Veterinary Medicines Authority**

**PERMIT TO ALLOW EMERGENCY USE OF A REGISTERED AGVET CHEMICAL
PRODUCT FOR CONTROL OF FALL ARMYWORM IN VARIOUS CROPS**

PERMIT NUMBER – PER89353

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 5 MAY 2020 TO 31 MAY 2023

Permit Holder:

HORTICULTURE INNOVATION AUSTRALIA LTD
Level 7, 141 Walker Street
NORTH SYDNEY NSW 2060

Persons who can use the product under this permit:

Persons generally.

CONDITIONS OF USE

Products to be used:

ALTACOR HORT INSECTICIDE (APVMA No. 69957)

Containing: 350 g/Kg CHLORANTRANILIPROLE as the only active constituent.

CORAGEN INSECTICIDE (APVMA No. 61519)

Containing: 200 g/L CHLORANTRANILIPROLE as the only active constituent.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The advisory buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas. DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are surface temperature inversion conditions present at the application site during the time of application. These conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers').

Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones for natural aquatic areas	
		WG formulation ALTACOR HORT INSECTICIDE	SC formulation CORAGEN INSECTICIDE
Up to maximum permit rate	0.5 m or lower	Not required.	Not required.
	1.0 m or lower	20 metres	25 metres

DO NOT apply by a vertical sprayer unless the following requirements are met:

- Spray is not directed above the target canopy.
- The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.
- For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers').

Buffer zones for vertical sprayers

Type of target canopy and dilute water rate	Mandatory downwind buffer zones for natural aquatic areas	
	WG formulation ALTACOR HORT INSECTICIDE	SC formulation CORAGEN INSECTICIDE
2 metres tall and shorter, maximum dilute water rate of 1000 L/ha	10 metres	10 metres
Taller than 2 metres (not fully-foliated), maximum dilute water rate of 1700 L/ha	25 metres	30 metres
Taller than 2 metres (fully-foliated), maximum dilute water rate of 1700 L/ha	20 metres	20 metres

Directions for Use:

Crop	Pest	Rate
<i>Rubus</i> spp. (field only)	Fall Armyworm (<i>Spodoptera frugiperda</i>)	<u>350 g/kg product</u> 18 g/100 L + 15 g a.i./100 L of non-ionic surfactant/wetting agent
Tree nuts (except almonds)		<u>200 g/L product</u> 30 mL/100 L + 15 g a.i./100 L of non-ionic surfactant/wetting agent
Strawberries (field and protected)		350 g/kg product 6 g/100 L or 60 g/ha + 15 g a.i./100 L of non-ionic surfactant/wetting agent
Parsley (field and protected)		200 g/L product 10 mL/100 L or 100 mL/ha + 15 g a.i./100 L of non-ionic surfactant/wetting agent
Root and tuber vegetables (except potatoes) (field only)		

Critical Use Comments:

- Regularly scout crops to monitor for eggs and larvae. Treat when pests appear, targeting eggs at hatch or small larvae (prior to third instar stage) before the pest becomes entrenched.
- Apply by boom spray or air-blast sprayer or equivalent.
- DO NOT apply more than 3 applications per crop.
- DO NOT apply more than 2 consecutive sprays per crop, with a minimum interval of 7 days.
- Further treatments should be made with a alternative mode of action insecticide.
- DO NOT apply less than 7 – 14 days after the initial treatment.
- Use enough water to ensure thorough coverage of the crop.
- Use a non-ionic surfactant/wetting agent at 15 g active/100 L, (e.g. Agral 600 @25 ml/100 L).
- DO NOT use BS1000 or Activator as it may cause phytotoxicity.
- Refer to the Insecticide Resistance Management (IRM) on the product label to prevent or delay the development of insecticide resistance.
- Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

Withholding Period:

Parsley, Rubus spp. and Root and tuber vegetables (except potato):

Harvest: Do not harvest for 3 days after application.

Strawberries:

Harvest: Do not harvest for 1 day after application.

Tree nuts (except almonds):

Harvest: Do not harvest for 10 days after application.

All crops:

Grazing: Do not graze or cut for stock food.

Jurisdiction:

All States and Territories, except VIC.

Note: Victoria is not included in this permit, as their Control-of-Use legislation means a permit is not required to legalise this off-label use in that State.

Additional Conditions:

This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label.

To Avoid Crop Damage

The sensitivity of the crop to be treated under this permit has not been fully evaluated. It is advisable to only treat a small area to ascertain the reaction before treating the whole crop.

Export of treated produce

Maximum Residue Limits (MRLs) and Temporary Maximum Residue Limits (TMRLs) have been established to allow treated produce to be used for human and animal consumption. An MRL has been established for CHLORANTRANILIPROLE. MRLs can be found in the *Agricultural and Veterinary Chemicals Code Instrument No. 4 (MRL Standard)*. MRLs apply only to produce marketed and consumed in Australia. If treated produce is to be exported, residues must not exceed the limits/tolerances of the importing country.

Issued by the Australian Pesticides and Veterinary Medicines Authority