TB Skin Test Reporting Form	Name
for Healthcare Professional	Date of Birth
Students	APID
	College

Date TB test placed	
Date TB test read	
Results in mm. of induration (hard, dense, and raised formation) and, if induration is present, measuring the diameter of induration transverse (perpendicular) to the long axis of the forearm. Erythema or redness of the skin should not be considered when reading a TB skin test result.)r
Interpretation of result by Centers for Disease Control and Prevention Guidelines (CDC)- See below: Negative Positive	
Signature of Physician or Registered Nurse	
Printed name	

CDC Classification of the Tuberculin Skin Test Reaction

An induration of 5 or more millimeters is	An induration of 10 or more millimeters is	An induration of 15 or more
considered positive in	considered positive in	millimeters is considered positive
-HIV-infected persons -A recent contact of a person with TB disease	-Recent immigrants (< 5 years) from high- prevalence countries -Injection drug users	in any person, including persons with no known risk factors for TB. However, targeted skin testing programs should only be conducted among high-risk groups.
-Persons with fibrotic changes on chest radiograph consistent with prior TB	-Residents and employees of high-risk congregate settings	
-Patients with organ transplants	-Mycobacteriology laboratory personnel	
other reasons (e.g., taking the equivalent	1 ~	
of >15 mg/day of prednisone for 1 month or longer, taking TNF-a antagonists)	-Children < 4 years of age	
	- Infants, children, and adolescents exposed to adults in high-risk categories	

• Please note-All incoming medical and nursing students must have a two-step tuberculin skin test, even if they have been tested annually prior to entering their medical/nursing program. The second step tuberculin test must be read 1-3 weeks after the first test.

Send results to:
Office of the University Physician
463 E. Circle Dr, Room 123
East Lansing, MI 48824-1037
Or fax to: 517.355.0332