

Bebtelovimab Monoclonal Therapy EUA Guidance for COVID-19

ELIGIBILITY

Treatment of mild to moderate COVID-19 disease in patients:

- ✓ Positive SARS-COV 2 test
- ✓ Within **7 days** of symptom onset
- ✓ Age ≥ 12 years old
- ✓ Weight ≥ 40kg
- ✓ O2 sats ≥ 94% and not requiring increased supplemental oxygen in baseline rate due to COVID
- ✓ for whom alternative COVID-19 treatment options approved or authorized by FDA are not accessible or clinically appropriate

Have **ANY** of the following risk factors for progression to severe disease:

- BMI > 25 (or if age 12-17 BMI ≥ 85th percentile for age group)
- Chronic Kidney Disease
- Diabetes Mellitus
- Immunosuppressive disease or immunosuppressive treatment
- Age ≥ 65 years
- Cardiovascular disease
- Hypertension
- Pregnancy
- Sickle cell disease
- Congenital or acquired heart disease
- Medical related technological dependence (i.e. tracheostomy)
- Asthma, COPD, or other chronic respiratory disease

[Bebtelovimab Health Care Provider Fact Sheet 03252022 \(fda.gov\)](#)

DOSE

- Bebtelovimab 175mg as a single IV Push over at least 30 seconds
- Flush the injection line with 10 mL of 0.9% Sodium Chloride to ensure delivery of the entire dose

HOW TO ORDER

- Educate patient and obtain verbal consent
- Search in Epic under “**Monoclonal therapy**”
- Choose the following panel:**
 - Treatment of Mild-Moderate COVID-19**
- The screening questions are built into the order set
- Send message to IC provider pool with patient name MRN (via telephone communication tool)
- The nurses will contact patient to schedule appointment
- Patient will be assessed upon arrival by urgent care provider and an order will be placed if patient is still meeting criteria