

Department of
Respiratory & Critical Care Medicine


Idiopathic Pulmonary Fibrosis (IPF)



Idiopathic Pulmonary Fibrosis

What is Idiopathic Pulmonary Fibrosis?

Idiopathic pulmonary fibrosis (IPF) is a progressive lung condition which causes scarring of the tissues (fibrosis) around the air sacs in the lungs. This scarring reduces the efficiency of oxygen and carbon dioxide exchange within the lungs and causes difficulties in breathing.



Who is Likely to Get IPF?

The exact cause of IPF is unknown. However, the risk of IPF is greater for:

- People who smoke or used to smoke
- People who are above the age of 50 years old
- Members of the same family (occasionally)

What are the Symptoms of IPF?

In the early stages of IPF, patients may experience mild or no symptoms. However, symptoms may worsen with increased scarring of the lungs.

Symptoms of IPF are not unique to this disease. While the symptoms may be different for each IPF patient, common symptoms associated with IPF include:

- Breathlessness, particularly when climbing up the stairs or walking uphill
- Persistent dry cough
- Tiredness

How is IPF Diagnosed?

Your doctor may ask you about your existing medical conditions, medications and occupation. He or she may also check for abnormal breathing sounds using a stethoscope.

If you are suspected of having IPF, the following tests may be done:

Blood Test



To exclude other possible causes of lung scarring

Breathing/ Lung Function Test



To measure how well your lungs are working

Computer Tomography (CT) Scan




To obtain a detailed imaging of your chest

Bronchoscopy



To obtain cell samples from your lungs

Lung Biopsy



To obtain tissue samples from your lungs

How is IPF Treated?


You can play your part through the following:



1. **Quit smoking** to prevent further damage to your lungs and ensure the effectiveness of your medications.
2. Attend **pulmonary rehabilitation** sessions to learn how to manage your symptoms and improve your physical strength.
3. Stay **active and healthy** by working out, eating well, and maintaining social activities and hobbies.
4. **Get vaccinated** to prevent infections from affecting your lungs.
5. Seek help if you are facing emotional or financial difficulties. Speak to your doctor for referrals to **support services** (e.g. medical social workers).

How is IPF Treated?

Your doctor may prescribe the following treatments:



1. **Medications** that may help slow down lung damage:
 - Nintedanib
 - Pirfenidone
2. **Oxygen therapy** to be used when you are feeling breathless, or throughout the day, depending on your condition.
3. **Lung transplantation** to replace one or both of your lungs with a healthy one from a donor. If this is an option, you will be referred to a transplant centre for evaluation.

Palliative Care and Advanced Care Planning



Palliative care is care for patients with moderate to severe symptoms that significantly affect their well-being.

Palliative care aims to improve a patient's quality of life by managing their symptoms such as breathlessness, cough and fatigue. This will allow people to continue leading meaningful lives and it also provides support to family members and caregivers.

Depending on your condition, your doctor may refer you to a multi-disciplinary team that specialises in providing palliative care.

Advanced Care Planning (ACP) involves discussing your personal values, beliefs, and care preferences with your family and healthcare professionals. ACP allows you to give instructions regarding the type of treatment you may or may not want, should you face a critical illness. It would relieve your family members from making difficult decisions for you, should you encounter a medical emergency.



To find out more about ACP, speak to your doctor or scan the QR code at the back of this booklet.

Brought to you by the Department of Respiratory & Critical Care Medicine, with inputs from Department of Pharmacy.

This information is updated and accurate as of July 2020.

This information is for educational purposes only and should not be used as a substitute for the professional medical advice or treatment.

Clinics 4A & 4B
TTSH Medical Centre, Level 4
Contact:
6357 7000 (Central Hotline)



Scan the QR Code with your smart phone to
access the information online or visit
<http://bit.ly/TTSHHealth-Library>

Was this information helpful?
Please feel free to email us if you have
any feedback regarding what you have
just read at
patienteducation@ttsh.com.sg



© Tan Tock Seng Hospital, Singapore 2021. All rights reserved. All information correct as of May 2021. No part of this document may be reproduced, copied, reverse compiled, adapted, distributed, commercially exploited, displayed or stored in a database, retrieval system or transmitted in any form without prior permission of Tan Tock Seng Hospital. All information and material found in this document are for purposes of information only and are not meant to substitute any advice provided by your own physician or other medical professionals.