

What will happen to my vision?

In most cases, vision starts to improve within one or two weeks of treatment. In severe cases, there may be a residual impairment of colour and brightness.

Will it happen again?

There is a possibility that optic neuritis will recur, or affect the other eye. If you experience similar symptoms in future, you should see your ophthalmologist or proceed to the Emergency Department immediately.



Clinic Appointments

Tel: (65) 6357 7000

Email: contact@ttsh.com.sg

Website: www.ttsh.com.sg

LASIK Enquiries

Tel: (65) 6357 2255

Email: lasik@ttsh.com.sg

Website: www.ttshlasik.com.sg



Scan the QR Code with your smart phone
to access the information online or visit
<http://bit.ly/TTSHHealth-Library>

Was this information helpful?
Please feel free to email us if you
have any feedback regarding what
you have just read at
patienteducation@ttsh.com.sg

Disclaimer

This is a general guide. If in doubt, please consult your doctor. Information is subject to revision without notice. The contents of this leaflet are not to be reproduced in any form without the prior permission of NHG Eye Institute.

DEPARTMENT OF OPHTHALMOLOGY

Optic Neuritis



What is Optic Neuritis?

Optic neuritis is a condition affecting the optic nerve. The optic nerve carries information from the eye to the brain, thus enabling us to see. In optic neuritis, there is inflammation of the optic nerve or the coverings of the nerve, causing the vision to be impaired.



Illustration of the visual pathway (view from side of the head).




Illustration of the visual pathway (view from top of the head).

The commonest symptoms are:

1. Poor vision which worsens over the first week
2. Decrease in colour vision
3. Decrease in light brightness and contrast
4. There may be pain, usually behind the eye, worsened by moving your eye in different directions.

What causes it?

There are many causes of optic neuritis, including:

1. **An overactive immune response** in your own body
 - This is the commonest cause.
 - The covering or myelin of the nerve is attacked by the immune system, and becomes inflamed, leading to "demyelination".
2. Rare conditions such as **multiple sclerosis** or **neuromyelitis optica**, which affect the brain and spinal cord as well.
3. **Other autoimmune conditions** which affects other parts of the body such as Sjogren's disease.
4. Certain types of **infections**.

Your doctor is likely to order a scan of your brain and nerves (usually an MRI scan) to confirm the diagnosis and to rule out any other diagnoses. Your doctor will also need to perform several other investigations ranging from blood tests to eye imaging tests, to confirm the diagnosis and that you are safe to receive treatment.

In cases where there is involvement of the brain, your doctor may refer you to the Neurology team for further investigations.

What is the treatment?

For the most common type of optic neuritis (demyelination), your vision is likely to recover over the next 4 to 6 weeks, even without treatment.

However, studies have shown that treatment with intravenous high-dose steroids (administered into the bloodstream) may quicken the recovery and reduce the short-term risk of multiple sclerosis.

As these steroids are given at a high dose, you will need to be monitored closely in the hospital. The steroids may be given more than once a day (in staggered doses), for a total of 3 to 5 days.

Potential side effects of steroids include:

1. Fever
2. High blood pressure
3. High blood sugar
4. Change in mood and insomnia
5. Bleeding from existing gut or stomach ulcers

To prevent these side effects, you will be given appropriate medication and monitored closely on the ward. Your doctor will discuss this in more detail with you.

