

Possible Risks of Surgery

The main risk of the surgery is to the recurrent laryngeal nerve, which is attached to the voice box and runs just next to the thyroid gland. Your voice may change if the nerve is damaged during the surgery.

The other main risk of surgery occurs only if both thyroid lobes are removed, which will affect calcium levels in your body.

The long term survival rates for most thyroid cancers are good and majority of patients have normal life expectancy after recovery.

Cancer Care Management



Our Surgeons are highly trained to ensure removal of the cancer and affected thyroid gland precision, while ensuring maximum function of the recurrent laryngeal nerve and preservation of the parathyroid glands.

Our aim is the same as yours – to fight off the cancer and return you to an ideal quality of life.

Clinics 1B

TTSH Medical Centre, Level 1

Contact:

6357 7000 (Central Hotline)



Scan the QR Code with your smart phone
to access the information online or visit
<http://bit.ly/TTSHHealth-Library>

Was this information helpful?
Please feel free to email us if you
have any feedback regarding what
you have just read at
patienteducation@ttsh.com.sg

Department of
OTORHINOLARYNGOLOGY
(EAR, NOSE & THROAT)

Thyroid Cancer



Overview

The diagnosis of cancer in a patient is often accompanied by disbelief and feelings of helplessness. The feeling of helplessness may be partly due to a lack of knowledge and awareness of the many treatment options available.

This series of brochures will provide you information on common head and neck cancers, focusing on the following:

- Common causes and symptoms
- Investigations required
- Treatment options available

For more information about the condition and treatment options, please consult a head and neck cancer specialist.

Cancer Patient Services

The Department of Otorhinolaryngology (Ear, Nose, Throat) in Tan Tock Seng Hospital offers the following services in relation to head and neck cancers:

1. Diagnosis of the cancer
2. Appropriate staging of the cancer
3. Treatment – every cancer patient is discussed with a multi-disciplinary team to identify the most ideal evidence-based treatment
4. Long-term follow up and monitoring to ensure the cancer does not return

Thyroid Cancer

Thyroid cancer is a cancer that starts in the thyroid gland. The thyroid gland sits in the lower middle portion of the neck and consists of a right and left lobe.

Various cancers may arise in the thyroid gland, the most common of which is papillary thyroid cancer.

Main Causes

- Family history of thyroid cancer
- Previous exposure to radiation to the neck

However, quite often patients with thyroid cancer may have none of these risk factors.

Common Symptoms


- Noticeable lump in the neck.
- Hoarseness of voice
- Difficulty swallowing
- Enlarged neck lymph glands

Examination and Tests

After the examination, your doctor will arrange for two tests:

1. Blood test to look at your thyroid function
2. Ultrasound of the thyroid

If there are any suspicious features, your doctor will arrange a **needle test** to determine if the lump is cancerous.



Treatment

In general, most papillary thyroid cancers are curable with surgery, although additional treatment may be required.

Based on the results of the ultrasound and needle test, your doctor will explain the type of thyroid cancer you have, how it will affect you and whether surgery is recommended.

1. Surgery

During the surgery, your surgeon may remove either the thyroid lobe with the tumour or both thyroid lobes. The extent of surgery depends on several factors, such as your age and the size of the tumour.

2. Radioactive Iodine Therapy

This once-off treatment is usually recommended for aggressive cancers.

You will be asked to ingest radioactive iodine in liquid form, which targets and kills cancerous thyroid cells in any part of the body. There is little effect on the rest of the body.