

Colonoscopy: Gastroenterology Procedure Unit

Screening Saves Lives

Colorectal Cancer is the second leading cancer killer in the United States. If you are 45 or older or have a family history of cancer, getting a colorectal screening can save your life.



- Colorectal Cancer often starts from a non-cancerous growth called a polyp that is in the rectum or colon.
- A colonoscopy can find **polyps** so that they can be removed before they turn into cancer.
- A colonoscopy can also find colorectal cancer early when the chances of being cured is good.



Colon Polyp

Preparing For Your Exam: What to Bring

- An adult (18 or older) to drive you home. You can take a cab, medical transport or bus only if an adult is riding with you after your visit.
- All medications you take, even over the counter medications
- Any vitamins and supplements you take
- Your insurance card

Medications

Call your doctor for medication instructions at least **one week before** your exam if you:

- Take diabetes medication
- Have a heart valve replacement
- Take Plavix, Brilinta or Aggrenox, or any other antiplatelet medications
- Take Coumadin (Warfarin), Xarelto, Pradaxa, Eliquis, Savaysa or other blood thinning medications
- Continue taking Aspirin unless told not to

Suflave™ Bowel Prep

5 Days Before Your Exam

Fill your bowel prep prescription and review these instructions.

You will get 2 bottles and 2 flavor packets.

The colon must be cleared of all solid mater so that the doctor can see clearly.

If your colon is not clean your exam may be cancelled.



3 Days Before Your Exam

- **Do not** eat any raw fruits, raw vegetables, seeds or nuts and no dairy.
- Stop iron and fiber supplements
- Mix and refrigerate bowel prep solution
- Plan for your ride home

Colonoscopy: Preparing for Your Exam

2 Days Before Your Exam

Do not eat any solid food. Begin clear liquid diet in the morning.

- **Examples of clear liquids are:** clear broth (chicken or beef), water, apple juice, Gatorade, popsicles and coffee or tea (no milk or creamer). You can have hard candy.
- Do not have any red or purple liquids
- Drink **at least 8 glasses of water** or clear liquid to stay hydrated.

The Day Before Your Exam

Do not eat any solid food. Start drinking bowel prep.

- Pour 1 flavor packet into 1 bottle of Suflave™. Fill the bottle up to the fill line with lukewarm water. Put the cap on the bottle and shake until the powder is mixed well. For the best taste, keep the mixture in the fridge for 1 hour before drinking.
- Drink 8 ounces of the mixture every 15 minutes until the bottle is empty.
- Drink at least 2 glasses of water during the evening.
- If you throw up, wait 30 minutes and begin drinking $\frac{3}{4}$ of a cup (6 ounces) every 15 minutes until half of the bottle is finished.

Tips

- Drink the dose quickly.
- Rinse your mouth with water or mouthwash.
- Try drinking with a straw.
- Suck on hard candy.

5 to 8 Hours Before Your Exam

- Mix the second bottle of Suflave™.
- Drink 8 ounces of the mixture every 15 minutes until the bottle is empty.
- Drink at least 2 glasses of water during the morning.
- Take your doctor approved medication.
- You can take aspirin, blood pressure medication, prednisone or other steroids.
- Keep a clear liquid diet.
- Your stool should be a clear yellow liquid.

2 Hours Before Your Exam

Do not eat or drink anything or your exam may be cancelled.

To make, change, or cancel an appointment, call (773) 702-6767.

Arrive to your exam 1 hour early.

- Hyde Park Location:** Center for Care and Discovery (CCD) 7th Floor
5700 South Maryland Ave, Chicago, IL 60637 **(773) 702-6767**
- Tinley Park Location:** Ingalls Family Care Center, Entrance C
6701 West 159th Street Tinley Park, IL 60477 **(708) 429-0222**
- River East Location:** UChicago Medicine River East
355 East Grand Ave, Chicago, IL 60611 **(773) 795-4662**
- Orland Park Location:** 14290 South Lagrange Road,
Orland Park, IL 60462 **(773) 834-9445**