Button Gastrostomy Tube (G-Tube) Care Tips Low Profile Button G-Tube There are 2 brands with similar Low profile Button G-Tubes are sized so that: features: MIC-KEY and AMT Access port Access port connects to extension set lavs flush (flat) on the skin Balloon port to fill • Balloon rests in balloon with water the stomach holding the Balloon button in place Positioning After Surgery (For babies under 6 months) Before your baby is positioned on their stomach, you must stop and disconnect the feeding and extension set. • After 4 days: Baby may sit up or lay against your chest. • For weeks 2 to 3: Baby may lay on tummy with a roll under upper chest for support. At 4 weeks: Baby may lay on tummy without roll. Feeding with a G-Tube Before you go home you will be given a home feeding plan. **Extension Set** Your surgery team will show you: How to give your child's feeding • How to flush the G-Tube Button G-Tubes must have an extension set to connect to the feeding tubing. To connect the extension set: Match the black lines on the extension set and the access port AFR • Push extension set down and turn to the right to lock in place Line Up Black Marks To disconnect the extension set: • Turn to the left and lift extension set from access port ENFIT Always disconnect the extension set when not in use Connection The extension set connects to the feeding tubing with an "ENFIT" connection. The 'ENFIT' connection screws together and is used only for G-Tube feedings Your feeding tubing and syringes will have this special connection Flushing the G-Tube Flush the G-tube with 5ml of water before and after each feeding and medication.

- If not using the G-tube every day, flush at least one time a day with 5 to 10ml of water to
 prevent clogging.
- Remove extension set from the Button G-Tube after flushing.



AT THE FOREFRONT OF K MEDICINE" UChicago Medicine Comer Children's

Health Literacy and Plain Language Translation by Diversity, Equityand Inclusion Department 10-11-2021Page 1 of 2

Button Gastrostomy Tube (G-Tube) Care Tips

General Care		
Site Care:	Bathing:	Securing:
 Clean the G-Tube site with gentle soap and water 2 times a day and as needed to keep the site from crusting. Turn the G-Tube several times a day to keep it from staying in the same spot. 	 Sponge bathe for the 1st month after G-Tube is placed. When given the OK by APN, you may give your child a regular bath in a tub. 	 Secure the G-Tube under clothing if needed. Use one piece clothing or clothing that helps keep child's hands from pulling on the G-Tube.

G-Tube Leakage

A small amount of leaking around the G-Tube is normal and can be managed with daily cleaning. If feeding or medications are leaking out around the tube, or if the surrounding skin becomes red and irritated, call the Pediatric Surgery APN.

You can check and refill the water in the balloon if your care team has instructed you how to do it. Refiling the balloon to the correct amount of water may lessen leaking.

Granulation Tissue

Granulation tissue is an "overgrowth" of tissue around the G-Tube. It develops when the body tries to repair itself at the G-Tube site.

If it bleeds or there is a large amount of tissue, call the Pediatric Surgery APN for treatment options.



Follow-up Appointments

- 2-week follow-up: Your first visit after G-Tube surgery may be an online or virtual visit.
- **3-month follow-up:** This will be an in-clinic visit to check the G-Tube site and do the first G-Tube change.
- Follow-up visits will then be every 3 months in the clinic for G-Tube changes. Your surgery team may also teach you how to do this at home. If you are changing the G-Tube at home, your child should be seen in the clinic for a follow-up once a year.

When To Call Pediatric Surgery With Concerns (773) 702-6169

- Changes at G-Tube site: redness,
 Fe warmth, a lot of drainage or leaking,
 Vabad odor, pain or skin breakdown
 If y
- Fever over 101.5 degrees
 - **Vomiting** (throwing up) or Diarrhea (watery stools)
 - If you are not able to flush or use the G-Tube

If Your Child's G-Tube Comes Out

This is a serious problem that needs medical care right away because the G-Tube opening can close down quickly.

- Cover the site with a clean dry gauze or Band-Aid
- Get medical help right away
 - During the day **Monday to Friday**: Call the Pediatric Surgery APN to get an urgent clinic appointment at (773) 702-6169
 - **During the night or on weekends:** Go to the Emergency Room for a replacement



UChicago Medicine Comer Children's

Health Literacy and Plain Language Translation by Diversity, Equityand Inclusion Department 10-11-2021Page 2 of 2